Running title: Effect of circSEC24A on the progression of HCC

CircSEC24A promotes tumor progression through sequestering miR-455-3p in hepatocellular carcinoma

Yulin Liao¹, Dongming Yang², Huaichao Luo¹, Guishu Yang³, Xiaoxia Wen⁴, Kaijiong Zhang¹, Chang Liu¹, Ruiling Zu¹, Lichun Wu¹, Zuo Wang¹, Dongsheng Wang^{1,*}

¹Department of Clinical Laboratory, Sichuan Cancer Hospital and Institute, Sichuan Cancer Center, School of Medicine, University of Electronic Science and Technology of China, Chengdu, Sichuan, China; ²Department of Clinical Laboratory, The Fifth Affiliated Hospital of Sun Yat-sen University, Zhuhai, Guangdong, China; ³College of Clinical Medicine, Southwest Medical University, Luzhou, Sichuan, China; ⁴Clinical College, Chengdu University of Traditional Chinese Medicine, Chengdu, Sichuan, China

*Correspondence: dswang08@163.com

Received March 5, 2021 / Accepted May 27, 2021

Hepatocellular carcinoma (HCC) ranks third in the cause of death due to cancer. Circular RNA circSEC24 Homolog A (circSEC24A) has been uncovered to be upregulated in liver cancer. However, the function of circSEC24A in HCC is indistinct.

We analyzed the microarray datasets GSE78520 and GSE94508 to search for differentially expressed circRNAs associated with HCC. Expression of circSEC24A, microRNA (miR)-455-3p, and protein phosphatase, Mg²⁺/Mn²⁺ dependent 1F (PPM1F) mRNA was detected by quantitative real-time polymerase chain reaction (RT-qPCR). Loss-of-function experiments were conducted to validate the biological function of circSEC24A in HCC cells in vitro and in vivo. Protein levels were evaluated by western blotting and immunohistochemistry (IHC). The relationship between circSEC24A or PPM1F and miR-455-3p was verified by a dual-luciferase reporter and/or RNA immunoprecipitation (RIP) assays.

circSEC24A was overexpressed in HCC. circSEC24A silencing decreased xenograft tumor growth in vivo and repressed proliferation, metastasis, invasion, epithelial-to-mesenchymal transition (EMT), induced cell cycle arrest, and apoptosis of HCC cells in vitro. circSEC24A acted as a molecular sponge to sequester miR-455-3p, resulting in elevating the expression of PPM1F. miR-455-3p inhibitor reversed the suppressive impact of circSEC24A silencing on malignant behaviors of HCC cells. PPM1F overexpression offsets the inhibitory effect of miR-455-3p mimic on malignant behaviors of HCC cells.

circSEC24A sponged miR-455-3p to elevate the PPM1F expression, resulting in accelerating malignant behaviors of HCC cells. The study provided a potential therapeutic target for patients with HCC.

Key words: HCC; circSEC24A; miR-455-3p; PPM1F

46
47
48 Hepatocellular carcinoma (HCC) ranks third in the cause of death due to cancer [1]. The occurrence
49 of HCC is closely related to chronic hepatitis, especially chronic viral hepatitis infection (hepatitis
50 B or C) [2]. Most HCC patients are diagnosed at an advanced stage and are not suitable for surgical
51 treatment [3]. Also, patients with advanced HCC have an extremely poor prognosis [4]. Thus,
52 studying the underlying molecular mechanisms of HCC development is essential to provide
53 guidance for the exploitation of HCC treatment tactics.

54 Circular RNAs (circRNAs) are endogenous RNA molecules with unique circular structures. They
55 are derived from pre-mRNA through back-splicing and have the ability to resist RNase R digestion
56 [5]. CircRNAs are emerging as vital regulators of gene expression and cellular functions [6].
57 Mounting reports have uncovered that circRNAs are implicated in the development of HCC [7]. For
58 example, circRNA circLRIG3 drove tumor progression through regulation of the EZH2/STAT3
59 signaling [8]. Moreover, circRNA circSORE sustained cell chemoresistance through activating the
60 Wnt/ β -catenin pathway in HCC [9]. Also, circRNA circBACH1 repressed p27 expression by
61 interacting with HuR, resulting in accelerating cell growth in HCC [10]. Hsa_circ_0003528
62 (circSEC24A), located at chr5: 134032815-134044578, is generated by the SEC24 Homolog A,
63 COPII Coat Complex Component (SEC24A) gene. Huang et al. discovered that circSEC24A
64 contributed to the polarization of tuberculosis-related macrophages [11]. In addition, circSEC24A
65 had been revealed to be upregulated in liver cancer (GSE78520 and GSE94508). However, the
66 function of circSEC24A in cancer and HCC has not been validated.

67 Studies have demonstrated that circRNAs can sequester microRNAs (miRs) by miR response
68 elements, thus regulating downstream targets of miRs [12, 13]. MiR-455-3p is encoded by the
69 COL27A1 gene [14]. Abnormal miR-455-3p expression is related to a variety of human diseases,
70 such as osteoarthritis [15], pulmonary arterial hypertension [16], and cancer [17]. MiR-455-3p had
71 been disclosed as a tumor repressor in HCC [18, 19]. Nevertheless, the mechanism of miR-455-3p
72 dysregulation in HCC remains unclear. Members of the metal-dependent protein phosphatase (PPM)
73 family are negative regulators of cellular stress response pathways, such as HOG, JNK, and p38
74 MAPK signaling pathways [20]. Mg^{2+}/Mn^{2+} -dependent protein phosphatase 1F (PPM1F), a
75 serine/threonine protein phosphatase, belongs to the PPM family. PPM1F is implicated in cancer
76 cell motility and adhesion [21, 22]. Also, PPM1F plays dual regulatory functions in cancer cell

metastasis [23]. In gastric cancer, PPM1F had been unmasked as a suppressive factor [24], while PPM1F was proved as an oncogene in HCC [25, 26]. At present, the connection between miR-455-3p and PPM1F in HCC is unclear.

In the research, circSEC24A was confirmed to exert an oncogenic role in HCC. Moreover, circSEC24A elevated PPM1F expression by sequestering miR-455-3p, thus accelerating HCC cell proliferation, metastasis, invasion, and EMT.

Patients and methods

Clinical specimens and ethics approval. Paired HCC tissues and periphery (peri)-tumor tissues were collected from the Sichuan Cancer Hospital and Institute. All registered HCC patients had signed informed consents and had never suffered other tumors. The current research was ratified by the Ethics Committee of the Sichuan Cancer Hospital and Institute.

Cell culture. 293T cells and normal liver cells (THLE-2) were purchased from Mingzhou Biotechnology (Ningbo, China). HCC cells (Huh7, SK-Hep-1, Hep3B, and Bel-7405) were bought from Procell (Wuhan, China). These cells were cultured in Dulbecco's Modified Eagle Medium (Procell) (Huh7 and 293T cells), Minimum Essential Medium (Thermo Fisher, Carlsbad, CA, USA) (SK-Hep-1 and Hep3B cells), or Roswell Park Memorial Institute-1640 Medium (Thermo Fisher) (Bel-7405 and THLE-2 cells) supplemented with fetal bovine serum (FBS) (Thermo Fisher) and 1% Penicillin/Streptomycin (Thermo Fisher) under suitable conditions (95% relative humidity, 5% carbon dioxide, and 37 °C).

Oligonucleotides, plasmids, and transfection. Small interference (si) RNAs (si-circSEC24A#1, si-circSEC24A#2, and si-circSEC24A#3) that were used to silence endogenous circSEC24A expression, negative control for siRNAs (si-NC), short hairpin (sh) RNA against circSEC24A (sh-circSEC24A), negative control for shRNA (sh-NC), miR-455-3p inhibitor, negative control for miR-455-3p inhibitor (inhibitor NC), miR-455-3p mimic, and negative control for miR-455-3p mimic (mimic NC) were synthesized by RiboBio (Guangzhou, China). Overexpression plasmid of PPM1F (pcDNA-PPM1F) was constructed with the pcDNA vector (Thermo Fisher). Liposomal cocktails with oligonucleotides or/and plasmids were generated with Lipofectamine 3000 (Thermo Fisher). Transfected cells were harvested and used in subsequent experiments.

Quantitative real-time polymerase chain reaction (RT-qPCR). The TRIzol™ Plus RNA

107 Purification Kit (Thermo Fisher) was utilized for total RNA isolation. RNase R treatment was
108 carried out at 37 °C with 3 U/μg RNase R (BioVision, Milpitas, CA, USA). Reverse transcription
109 was performed using Prime Script™ RT reagent kit (TaKaRa, Dalian, China) or miRscript cDNA
110 synthesis kit (QIAGEN, Hilden, Germany). qPCR was set up with the SYBR Premix Ex Taq Kit
111 (Applied Biosystems, Foster City, CA, USA) in triplicate and the primers utilized in the present
112 research were exhibited in Table 1. U6 or GAPDH was used as an endogenous reference. Data was
113 figured with the $2^{-\Delta\Delta C_t}$ method.

114 **Cell counting kit-8 (CCK-8) assay.** Treated HCC cells (about 2×10^3) were cultured in 96-well
115 plates. After incubation with CCK-8 solution (10 μl, Dojindo, Kumamoto, Japan), the optical
116 density (OD) value was read with the spectrophotometer reader (Thermo Fisher) at a wavelength of
117 450 nm.

118 **Colony formation assay.** Treated HCC cells (approximately 2×10^3) were seeded into 6-well
119 plates. The colonies were stained with crystal violet solution (Sigma, St. Louis, MO, USA) (0.5%
120 crystal violet in 10% methanol and 90% water) and then counted with a microscope (Olympus,
121 Tokyo, Japan).

122 **Flow cytometry assay.** HCC cells in the logarithmic phase were harvested and fixed with cold 70%
123 ethanol. Subsequently, the cells were digested with 2 μg RNase A (Sigma,) and then labeled with 15
124 μg/ml propidium iodide (PI) (Beyotime, Jiangsu, China) for 15 min. The cell cycle distribution was
125 analyzed using the FACS Calibur flow cytometer (Becton Dickinson, San Jose, California, USA).
126 Cell apoptosis analysis was carried out according to the instructions of the Annexin V-FITC/PI
127 apoptosis detection kit (Sigma).

128 **5-ethynyl-2'-deoxyuridine (EDU) assay.** EDU assay was conducted with the Cell-Light EdU
129 Apollo 567 In Vitro Imaging Kit (RiboBio, Guangzhou, China) based on the manufacturer's
130 instructions. Images were captured by the microscope (Olympus). EDU-positive cells were labeled
131 with red fluorescence, while nuclei were labeled with DAPI (4,6-diamidino-2-phenylindole; blue).

132 **Transwell assay.** The metastasis and invasion of HCC cells were analyzed by transwell assay with
133 transwell chambers with (#354480, Costar, Cambridge, MA, USA) or without Matrigel. In short,
134 treated HCC cells (5×10^4) in a serum-free medium were seeded onto the apical chamber. 24 h after
135 seeding, the migrated and invaded cells were stained with 0.5% crystal violet (Sigma) and then
136 counted in 5 randomly selected fields using a microscope (Olympus).

137 **Western blotting.** Total protein was extracted using the RIPA buffer (Beyotime) and then isolated
 138 with 10% sodium dodecyl sulfate-polyacrylamide gel electrophoresis. After transferring to the
 139 polyvinylidene fluoride (PVDF) membrane (Millipore, Bedford, MA, USA) and sealing with 5%
 140 skimmed milk powder, the membranes were incubated with the appropriate primary antibodies
 141 against E-cadherin (1:600, ab1416), Vimentin (1:1000, ab8978), PPM1F (1:1000, ab200394), and
 142 GAPDH (1:2500, ab8245), followed by washing and incubating with the appropriate peroxidase
 143 secondary antibody. These antibodies were purchased from Abcam (Cambridge, MA, USA). Bands
 144 were detected using the Pierce™ ECL WB Substrate (Thermo Fisher) and visualized after exposure
 145 onto HyBlot CL Autoradiography Film (Denville Scientific, Metuchen, NJ USA).

146 **Dual-luciferase reporter assay.** The binding sites of circSEC24A and PPM1F 3' untranslated
 147 region (UTR) in miR-455-3p were predicted using the starBase database. The wild type sequences
 148 of circSEC24A and PPM1F 3'UTR and their mutant sequences were inserted into the
 149 pMIR-REPORT vector (Applied Biosystems) to construct a luciferase vector carrying circSEC24A
 150 wt, PPM1F 3'UTR wt, circSEC24A mut, or PPM1F 3'UTR mut, respectively. Following this, 293T
 151 cells were transfected with mimic NC or miR-455-3p mimic together with a luciferase vector,
 152 followed by analyzing luciferase activities using the Pierce™ Renilla-Firefly Luciferase Dual Assay
 153 Kit (Thermo Fisher) based on the manufacturer's instructions on an LD400 luminescence detector
 154 (Beckman Coulter, Brea, CA, USA).

155 **RNA immunoprecipitation (RIP) analysis.** The interaction between circSEC24A and miR-455-3p
 156 was verified with the Magna RIP kit (Millipore) in accordance with the instructions from the
 157 manufacturer. Abundance of circSEC24A and miR-455-3p in precipitated RNAs was assessed
 158 through carrying out RT-qPCR analysis.

159 **Xenograft assay.** 14 BALB/c nude mice (Vital River Laboratory, Beijing, China) were manipulated
 160 based on protocols approved by the Animal Care Committee of Sichuan Cancer Hospital and
 161 Institute. Briefly, Hep3B cells carrying lentivirus-mediated sh-circSEC24A or sh-NC were
 162 subcutaneously injected into nude mice. The size of tumor was measured once a week
 163 ($\text{Volume} = (\text{length} \times \text{width}^2)/2$). All mice were sacrificed after 35 days, and xenograft tumors were
 164 stripped and weighed, followed by embedding in paraffin for immunohistochemistry (IHC)
 165 staining.

166 **Statistical analysis.** Statistical analyses were conducted using SPSS Statistics software version 17.0

167 (SPSS, Chicago, IL, USA). For *in vivo* and *in vitro* experiments, the Student's *t*-test or one-way
168 analysis of variance was utilized to evaluate the difference between two or among multiple groups.
169 $P < 0.05$ was considered statistically significant. All data were presented as mean $\pm$ standard
170 deviation from at least three repeats. Each experiment was conducted in triplicate.

171

172 **Results**

173 **CircSEC24A was highly expressed in HCC.** To search for differentially expressed circRNAs
174 related to HCC, we analyzed microarray datasets (GSE78520 and GSE94508) concerning the
175 circRNA profiles in liver cancer tissues. We found that there were 5 overlapping circRNAs
176 (hsa_circ_0067934, hsa_circ_0003528, hsa_circ_0001955, hsa_circ_0008274, and
177 hsa_circ_0072088) in the top 50 upregulated circRNAs in the GSE78520 and GSE94508
178 microarray datasets (Figure 1A). Among the 5 overlapping circRNAs, the biological role of
179 hsa_circ_0067934 [27], hsa_circ_0001955 [28], hsa_circ_0008274 [29], and hsa_circ_0072088 [30]
180 in HCC had been verified. Thus, we choose hsa_circ_0003528 for further exploration.
181 Hsa_circ_0003528 (circSEC24A) is formed by reverse splicing of exon 14 to exon 18 (741 bp) of
182 the SEC24A gene (Figure 1B). To verify the expression tendency of circSEC24A in 40 paired HCC
183 tissues and peri-tumor tissues, RT-qPCR was performed. As presented in Figure 1C, circSEC24A
184 expression was observably elevated in HCC tissues in contrast to peri-tumor tissues. Furthermore,
185 the Kaplan-Meier curve showed that HCC patients with high circSEC24A expression had a shorter
186 survival time than those with low circSEC24A expression (Figure 1D). As expected, circSEC24A
187 was upregulated in HCC cells (Huh7, SK-Hep-1, Hep3B, and Bel-7405) than that in the THEL-2
188 cells (Figure 1E). Then, RNase R treatment was performed to validate the circular structure of
189 circSEC24A. RT-qPCR exhibited that linear SEC24A was digested after RNase R treatment, but
190 circular SEC24A did not change. The digestion resistance of circSEC24A to RNase R confirmed its
191 closed-loop structure (Figure 1F). These results indicated that high circSEC24A expression might
192 be associated with the progression of HCC.

193 **CircSEC24A accelerated HCC cell proliferation, metastasis, invasion, and EMT.** Based on the
194 upregulation of circSEC24A in HCC, we investigated the biological function of circSEC24A by
195 loss-of-function experiments. Three siRNAs (si-circSEC24A#1, si-circSEC24A#2, and
196 si-circSEC24A#3) against the reverse splice region of circSEC24A were designed and transfected

197 into HCC cells. The knockdown efficiencies of three siRNAs were exhibited in Fig. 2A. CCK-8 and
198 colony formation assays indicated that circSEC24A silencing constrained growth and colony
199 formation of Huh7 and Hep3B cells (Figures 2B-2D). Cell cycle analysis revealed that circSEC24A
200 inhibition elevated the proportion of Huh7 and Hep3B cells in the G0/G1 stage and decreased the
201 proportion of Huh7 and Hep3B cells in the S stage, illustrating circSEC24A knockdown induced
202 cell cycle arrest in Huh7 and Hep3B cells (Figures 2E, 2F). EDU assay exhibited that circSEC24A
203 inhibition decreased the proportion of EDU-positive Huh7 and Hep3B cells (Figures 2G, 2H). Cell
204 apoptosis analysis presented that the proportion of apoptotic Huh7 and Hep3B cells was increased
205 after circSEC24A silencing (Figures 2I, 2J). Transwell assay showed that the inhibition of
206 circSEC24A decreased the number of migrated and invaded Huh7 and Hep3B cells (Figures
207 2K-2N). Moreover, circSEC24A silencing elevated E-cadherin protein level and decreased
208 Vimentin protein level in Huh7 and Hep3B cells (Figures 2O, 2P). These results manifested that
209 circSEC24A facilitated HCC cell proliferation, metastasis, invasion, and EMT *in vitro*.

210 **CircSEC24A was validated as a miR-455-3p sponge.** Many studies have revealed that circRNAs
211 can function as miR sponges in tumor development. By analyzing the starBase database, we
212 discovered that miR-455-3p might interact with circSEC24A and the binding sites of circSEC24A
213 in miR-455-3p were presented in Figure 3A. Dual-luciferase reporter assay exhibited that the
214 luciferase activity of the circSEC24A wt reporter was repressed in miR-455-3p mimic-transfected
215 293T cells (Figure 3B). RIP assay presented that circSEC24A and miR-455-3p could be co-enriched
216 in the anti-Ago2 group compared to the IgG group (Figure 3C). As expected, there was a significant
217 downregulation of miR-455-3p in HCC tissues and cells (Figures 3D, 3E). Also, the expression of
218 circSEC24A was negatively correlated with miR-455-3p in HCC tissues (Figure 3F). Together,
219 these results manifested that circSEC24A served as a sponge for miR-455-3p.

220 **CircSEC24A promoted HCC cell proliferation, metastasis, invasion, and EMT via sponging**
221 **miR-455-3p.** As we previously proved that circSEC24A acted as a miR-455-3p decoy, we
222 hypothesized that circSEC24A inhibition might reduce HCC cell proliferation, metastasis, invasion,
223 and EMT via sponging miR-455-3p. The silence efficiency of miR-455-3p inhibitor in Huh7 and
224 Hep3B cells was verified by RT-qPCR (Figure 4A). Moreover, miR-455-3p inhibitor reversed the
225 upregulation of miR-455-3p in circSEC24A-silenced Huh7 and Hep3B cells (Figure 4B). Also,
226 miR-455-3p inhibitor offset circSEC24A knockdown-mediated repressive influence on growth,

227 colony formation, cell cycle progression, and proliferation of Huh7 and Hep3B cells (Figures
228 4C-4F). Furthermore, the elevation of the proportion of apoptotic Huh7 and Hep3B cells caused by
229 circSEC24A silencing was overturned after miR-455-3p inhibition (Figure 4G). Additionally,
230 miR-455-3p inhibitor counteracted the suppressive effect of circSEC24A knockdown on migration
231 and invasion of Huh7 and Hep3B cells (Figures 4H-4K). As expected, miR-455-3p inhibitor
232 reversed the upregulation of E-cadherin protein and downregulation of Vimentin protein in
233 circSEC24A-silenced Huh7 and Hep3B cells (Figures 4L, 4M). Collectively, these results illustrated
234 that circSEC24A accelerated proliferation, metastasis, invasion, and EMT of HCC cells via
235 sponging miR-455-3p.

236 **PPM1F acted as a miR-455-3p target.** Based on the competing endogenous RNA (ceRNA)
237 hypothesis, circRNA can relieve the suppressive influence of miR on its target by functioning as a
238 miR sponge. Thus, we identified the potential target of miR-455-3p with the starbase database. As
239 exhibited in Figure 5A, PPM1F might be a target of miR-455-3p. The luciferase activity of the
240 PPM1F 3'UTR wt reporter was impaired by miR-455-3p mimic compared with the control group,
241 but the luciferase activity of the PPM1F 3'UTR mut reporter did not change (Figure 5B). Results of
242 RT-qPCR and western blotting revealed that the levels of PPM1F mRNA and protein were
243 upregulated in HCC tissues compared to peri-tumor tissues (Figures 5C, 5D). Also, the level of
244 PPM1F protein was upregulated in HCC cells (Figure 5E). Furthermore, there was a negative
245 correlation between PPM1F mRNA and miR-455-3p expression in HCC tissues (Figure 5F). The
246 decrease in PPM1F expression in si-circSEC24A#1-transfected Huh7 and Hep3B cells was reversed
247 by miR-455-3p inhibitor (Figure 5G). Thus, these results indicated that circSEC24A regulated
248 PPM1F expression through serving as a miR-455-3p sponge.

249 **MiR-455-3p repressed proliferation, metastasis, invasion, and EMT of HCC cells by targeting**
250 **PPMF1.** To investigate whether miR-455-3p played its function through targeting PPMF1 in HCC
251 cells, rescue experiments were conducted by transfecting pcDNA-PPMF1 into Huh7 and Hep3B
252 cells carrying miR-455-3p mimic. The overexpression efficiencies of miR-455-3p mimic and
253 pcDNA-PPMF1 in Huh7 and Hep3B cells were verified by RT-qPCR and western blotting,
254 respectively (Figures 6A, 6B). The transfection of pcDNA-PPMF1 restored the downregulation of
255 PPMF1 protein in Huh7 and Hep3B cells carrying miR-455-3p mimic (Figure 6C). Moreover,
256 PPMF1 overexpression overturned the inhibitory effect of miR-455-3p mimic on growth, colony

257 formation, cell cycle progression, and proliferation of Huh7 and Hep3B cells (Figures 6D-6G).
258 Furthermore, miR-455-3p mimic induced cell apoptosis and repressed cell metastasis and invasion
259 in Huh7 and Hep3B cells, but these impacts mediated by miR-455-3p mimic was offset after
260 pcDNA-PPMF1 introduction (Figures 6H-6L). In addition, miR-455-3p mimic elevated the
261 E-cadherin protein level and decreased the Vimentin protein level in Huh7 and Hep3B cells, but
262 these trends caused by miR-455-3p mimic were reversed by forcing PPMF1 expression (Figures
263 6M, 6N). In summary, miR-455-3p repressed HCC cell proliferation, metastasis, invasion, and EMT
264 via targeting PPMF1.

265 **CircSEC24A knockdown repressed HCC growth *in vivo*.** To further validate the function of
266 circSEC24A in HCC progression, Hep3B cells carrying lentivirus-mediated sh-circSEC24A or
267 sh-NC were subcutaneously injected into BALB/c nude mice. The results showed that tumor
268 volume and weight in the sh-circSEC24A growth were smaller and lower compared with the sh-NC
269 group (Figures 7A, 7B). As expected, the expression of circSEC24A and PPMF1 mRNA was
270 downregulated in mice tumor tissues in the sh-circSEC24A group in contrast to the sh-NC group,
271 but miR-455-3p expression was upregulated (Figure 7C). IHC analysis displayed that PPM1F and
272 Vimentin protein levels were downregulated in mice tumor tissues in the sh-circSEC24A group, but
273 the E-cadherin protein level was upregulated (Figure 7D). Together, the downregulation of
274 circSEC24A curbed HCC growth *in vivo*.

275

276 Discussion

277 A series of studies have found that differential circRNA expression is related to the clinical
278 characteristics of cancer patients [31, 32]. Also, circRNAs provide a new direction for studying
279 tumor pathogenesis and potential therapeutic targets [33]. Herein, we demonstrated that
280 circSEC24A played an oncogenic role through sequestering miR-455-3p and upregulating PPM1F
281 in HCC, offering evidence to support circSEC24A as an underlying therapeutic target for HCC.

282 The host gene of circSEC24A is SEC24A. SEC24A, a cytosolic protein, plays a fundamental role in
283 coat protein complex II (COPII)-dependent endoplasmic reticulum-Golgi protein transport [34].
284 Furthermore, the intracellular transport of hepatitis B subviral envelope particles depends on the
285 COPII mechanism through the selective utilization of SEC24A and SEC23B [35]. Also, SEC24A
286 had been uncovered to play a promoting impact on gastric cancer cell invasion and metastasis [36].

287 A recent report revealed that circSEC24A facilitated mycobacterium tuberculosis-related
288 macrophage polarization via upregulating CTLA4. These data offered evidence for the function of
289 circSEC24A. In this study, circSEC24A was highly expressed in HCC. And HCC patients with high
290 circSEC24A expression had a shorter survival time than those with low circSEC24A expression,
291 manifesting that circSEC24A might be a potential biomarker for analyzing the prognosis of HCC
292 patients. Moreover, circSEC24A silencing constrained xenograft tumor growth and repressed HCC
293 cell proliferation, metastasis, invasion, and EMT *in vitro*. Thus, these data indicated that
294 circSEC24A exerted a promoting effect on tumor progression in HCC.

295 Given that circRNAs can interact with miRs via miR response elements. Through a series of
296 validation, circSEC24A was validated as a miR-455-3p decoy. Report of Lan et al. discovered that
297 miR-455-3p downregulation-mediated STK17B upregulation activated the AKT/GSK-3 β /Snail
298 signaling, thus promoting tumorigenesis and metastasis in HCC [18]. It was reported that
299 miR-455-3p played an antagonistic effect on liver Fibrosis by targeting HSF1 [37]. Moreover,
300 TUG1 sequestered miR-455-3p and elevated AMPK β 2 expression, resulting in promoting cell
301 metastasis and glycolysis through in HCC [19]. Herein, miR-455-3p downregulation overturned
302 circSEC24A downregulation-mediated impacts on malignant behaviors of HCC cells. Consequently,
303 we deduced that circSEC24A contributed to HCC cell malignancy via serving as a miR-455-3p
304 sponge.

305 PPM1F, a serine/threonine phosphatase, takes part in the progression of human cancers [38]. A
306 previous study revealed that miR-590-3p promoted invasion and proliferation through targeting
307 PPM1F in gastric cancer [24]. In addition, Jurmeister et al. exposed that miR-200c inhibited cell
308 metastasis through downregulating FHOD1 and PPM1F [39]. In HCC, circRNA circSLC3A2
309 promoted cell invasion and proliferation through adsorbing miR-490-3p and upregulating PPM1F
310 [26]. Also, SNHG8 contributed to metastasis, invasion, and proliferation through regulation of the
311 miR-149-5p/PPM1F pathway [25]. Consistent with previous studies [25, 26], PPM1F was highly
312 expressed in HCC. Forced miR-455-3p expression repressed cell malignant behaviors via
313 decreasing PPM1F expression in HCC. Importantly, circSEC24A regulated PPM1F expression via
314 adsorbing miR-455-3p. Thus, we concluded that circSEC24A promoted tumor growth by regulation
315 of the miR-455-3p/PPM1F pathway in HCC.

316 In sum, circSEC24A was upregulated in HCC. Upregulation of circSEC24A adsorbed miR-455-3p

317 and increased PPM1F expression, thereby promoting tumor growth in HCC. The study manifested
318 that circSEC24A was a promising target for HCC treatment.

319

320

321 References

- 322 [1] MILLER KD, NOGUEIRA L, MARIOTTO AB, ROWLAND JH, YABROFF KR et al.
323 Cancer treatment and survivorship statistics, 2019. *CA Cancer J Clin* 2019; 69: 363-385.
324 <https://doi.org/10.3322/caac.21565>
- 325 [2] ALTER MJ. Epidemiology of hepatitis C virus infection. *World J Gastroenterol* 2007; 13:
326 2436-2441.
- 327 [3] ESCUDIER B, WORDEN F, KUDO M. Sorafenib: key lessons from over 10 years of
328 experience. *Expert review of anticancer therapy* 2019; 19: 177-189.
329 <https://doi.org/10.1080/14737140.2019.1559058>
- 330 [4] DISTEFANO JK, DAVIS B. Diagnostic and Prognostic Potential of AKR1B10 in Human
331 Hepatocellular Carcinoma. *Cancers (Basel)* 2019; 11.
332 <https://doi.org/10.3390/cancers11040486>
- 333 [5] GAN H, LEI Y, YUAN N, TANG K, HAO W et al. Circular RNAs in depression:
334 Biogenesis, function, expression, and therapeutic potential. *Biomed Pharmacother* 2021;
335 137: 111244. <https://doi.org/10.1016/j.biopha.2021.111244>
- 336 [6] RAJAPPA A, BANERJEE S, SHARMA V, KHANDELIA P. Circular RNAs: Emerging
337 Role in Cancer Diagnostics and Therapeutics. *Front Mol Biosci* 2020; 7: 577938.
338 <https://doi.org/10.3389/fmolb.2020.577938>
- 339 [7] CHENG D, WANG J, DONG Z, LI X. Cancer-related circular RNA: diverse biological
340 functions. *Cancer Cell Int* 2021; 21: 11. <https://doi.org/10.1186/s12935-020-01703-z>
- 341 [8] SUN S, GAO J, ZHOU S, LI Y, WANG Y et al. A novel circular RNA circ-LRIG3
342 facilitates the malignant progression of hepatocellular carcinoma by modulating the
343 EZH2/STAT3 signaling. *J Exp Clin Cancer Res* 2020; 39: 252.
344 <https://doi.org/10.1186/s13046-020-01779-5>
- 345 [9] XU J, WAN Z, TANG M, LIN Z, JIANG S et al. N-methyladenosine-modified
346 CircRNA-SORE sustains sorafenib resistance in hepatocellular carcinoma by regulating
347 β -catenin signaling. *Mol Cancer* 2020; 19: 163. <https://doi.org/10.1186/s12943-020-01281-8>
- 348 [10] LIU B, YANG G, WANG X, LIU J, LU Z et al. CircBACH1 (hsa_circ_0061395) promotes
349 hepatocellular carcinoma growth by regulating p27 repression via HuR. *J Cell Physiol* 2020;
350 235: 6929-6941. <https://doi.org/10.1002/jcp.29589>
- 351 [11] HUANG Z, YAO F, LIU J, XU J, GUO Y et al. Up-regulation of circRNA-0003528
352 promotes mycobacterium tuberculosis associated macrophage polarization via
353 down-regulating miR-224-5p, miR-324-5p and miR-488-5p and up-regulating CTLA4.
354 *Aging (Albany NY)* 2020; 12: 25658-25672. <https://doi.org/10.18632/aging.104175>
- 355 [12] HAN TS, HUR K, CHO HS, BAN HS. Epigenetic Associations between lncRNA/circRNA
356 and miRNA in Hepatocellular Carcinoma. *Cancers (Basel)* 2020; 12.
357 <https://doi.org/10.3390/cancers12092622>
- 358 [13] TAY Y, RINN J, PANDOLFI PP. The multilayered complexity of ceRNA crosstalk and
359 competition. *Nature* 2014; 505: 344-352. <https://doi.org/10.1038/nature12986>

- [14] KUMAR S, REDDY PH. A New Discovery of MicroRNA-455-3p in Alzheimer's Disease. *J Alzheimers Dis* 2019; 72: S117-S130. <https://doi.org/10.3233/JAD-190583>
- [15] HU S, ZHAO X, MAO G, ZHANG Z, WEN X et al. MicroRNA-455-3p promotes TGF- β signaling and inhibits osteoarthritis development by directly targeting PAK2. *Exp Mol Med* 2019; 51: 1-13. <https://doi.org/10.1038/s12276-019-0322-3>
- [16] ZHOU C, CHEN Y, KANG W, LV H, FANG Z et al. Mir-455-3p-1 represses FGF7 expression to inhibit pulmonary arterial hypertension through inhibiting the RAS/ERK signaling pathway. *J Mol Cell Cardiol* 2019; 130: 23-35. <https://doi.org/10.1016/j.yjmcc.2019.03.002>
- [17] LIU A, ZHU J, WU G, CAO L, TAN Z et al. Antagonizing miR-455-3p inhibits chemoresistance and aggressiveness in esophageal squamous cell carcinoma. *Mol Cancer* 2017; 16: 106. <https://doi.org/10.1186/s12943-017-0669-9>
- [18] LAN Y, HAN J, WANG Y, WANG J, YANG G et al. STK17B promotes carcinogenesis and metastasis via AKT/GSK-3 β /Snail signaling in hepatocellular carcinoma. *Cell Death Dis* 2018; 9: 236. <https://doi.org/10.1038/s41419-018-0262-1>
- [19] LIN YH, WU MH, HUANG YH, YEH CT, CHENG ML et al. Taurine up-regulated gene 1 functions as a master regulator to coordinate glycolysis and metastasis in hepatocellular carcinoma. *Hepatology* 2018; 67: 188-203. <https://doi.org/10.1002/hep.29462>
- [20] HANADA M, NINOMIYA-TSUJI J, KOMAKI K, OHNISHI M, KATSURA K et al. Regulation of the TAK1 signaling pathway by protein phosphatase 2C. *J Biol Chem* 2001; 276: 5753-5759. <https://doi.org/10.1074/jbc.M007773200>
- [21] SUSILA A, CHAN H, LOH AXW, PHANG HQ, WONG ET et al. The POPX2 phosphatase regulates cancer cell motility and invasiveness. *Cell Cycle* 2010; 9: 179-187. <https://doi.org/10.4161/cc.9.1.10406>
- [22] SINGH P, GAN CS, GUO T, PHANG H-Q, SZE SK et al. Investigation of POPX2 phosphatase functions by comparative phosphoproteomic analysis. *Proteomics* 2011; 11: 2891-2900. <https://doi.org/10.1002/pmic.201100044>
- [23] ZHANG S, WENG T, CHERUBA E, GUO T, CHAN H et al. Phosphatase POPX2 Exhibits Dual Regulatory Functions in Cancer Metastasis. *J Proteome Res* 2017; 16: 698-711. <https://doi.org/10.1021/acs.jproteome.6b00748>
- [24] ZHANG J, JIN M, CHEN X, ZHANG R, HUANG Y et al. Loss of PPM1F expression predicts tumour recurrence and is negatively regulated by miR-590-3p in gastric cancer. *Cell Prolif* 2018; 51: e12444. <https://doi.org/10.1111/cpr.12444>
- [25] DONG J, TENG F, GUO W, YANG J, DING G et al. lncRNA SNHG8 Promotes the Tumorigenesis and Metastasis by Sponging miR-149-5p and Predicts Tumor Recurrence in Hepatocellular Carcinoma. *Cell Physiol Biochem* 2018; 51: 2262-2274. <https://doi.org/10.1159/000495871>
- [26] WANG H, CHEN W, JIN M, HOU L, CHEN X et al. CircSLC3A2 functions as an oncogenic factor in hepatocellular carcinoma by sponging miR-490-3p and regulating PPM1F expression. *Mol Cancer* 2018; 17: 165. <https://doi.org/10.1186/s12943-018-0909-7>
- [27] ZHU Q, LU G, LUO Z, GUI F, WU J et al. CircRNA circ_0067934 promotes tumor growth and metastasis in hepatocellular carcinoma through regulation of miR-1324/FZD5/Wnt/ β -catenin axis. *Biochem Biophys Res Commun* 2018; 497: 626-632. <https://doi.org/10.1016/j.bbrc.2018.02.119>

- [28] DING B, FAN W, LOU W. hsa_circ_0001955 Enhances Proliferation, Migration, and Invasion of HCC Cells through miR-145-5p/NRAS Axis. *Mol Ther Nucleic Acids* 2020; 22: 445-455. <https://doi.org/10.1016/j.omtn.2020.09.007>
- [29] KONG Q, FAN Q, MA X, LI J, MA R. CircRNA circUGGT2 Contributes to Hepatocellular Carcinoma Development via Regulation of the miR-526b-5p/RAB1A Axis. *Cancer Manag Res* 2020; 12: 10229-10241. <https://doi.org/10.2147/CMAR.S263985>
- [30] YANG X, LIU L, ZOU H, ZHENG Y-W, WANG K-P. circZFR promotes cell proliferation and migration by regulating miR-511/AKT1 axis in hepatocellular carcinoma. *Dig Liver Dis* 2019; 51: 1446-1455. <https://doi.org/10.1016/j.dld.2019.04.012>
- [31] WANG M, YU F, LI P. Circular RNAs: Characteristics, Function and Clinical Significance in Hepatocellular Carcinoma. *Cancers (Basel)* 2018; 10: 258. <https://doi.org/10.3390/cancers10080258>
- [32] JIANG YL, SHANG MM, DONG SZ, CHANG YC. Abnormally expressed circular RNAs as novel non-invasive biomarkers for hepatocellular carcinoma: A meta-analysis. *World J Gastrointest Oncol* 2019; 11: 909-924. <https://doi.org/10.4251/wjgo.v11.i10.909>
- [33] BEILERLI A, GAREEV I, BEYLERLI O, YANG G, PAVLOV V et al. Circular RNAs as biomarkers and therapeutic targets in cancer. *Semin Cancer Biol* 2021; S1044-579X(21)00004-3. <https://doi.org/10.1016/j.semcancer.2020.12.026>
- [34] SHARPE LJ, LUU W, BROWN AJ. Akt phosphorylates Sec24: new clues into the regulation of ER-to-Golgi trafficking. *Traffic* 2011; 12: 19-27. <https://doi.org/10.1111/j.1600-0854.2010.01133.x>
- [35] ZEYEN L, DÖRING T, STIELER JT, PRANGE R. Hepatitis B subviral envelope particles use the COPII machinery for intracellular transport via selective exploitation of Sec24A and Sec23B. *Cell Microbiol* 2020; 22: e13181. <https://doi.org/10.1111/cmi.13181>
- [36] LU M, WANG W, ZHANG S, LI Y. SEC24A stimulates oncogenicity of human gastric cancer cells. *Int J Clin Exp Pathol* 2018; 11: 4044-4051.
- [37] WEI S, WANG Q, ZHOU H, QIU J, LI C et al. miR-455-3p Alleviates Hepatic Stellate Cell Activation and Liver Fibrosis by Suppressing HSF1 Expression. *Mol Ther Nucleic Acids* 2019; 16: 758-769. <https://doi.org/10.1016/j.omtn.2019.05.001>
- [38] KIM PR, ZHANG S, RAHMAT MB, KOH C-G. Partners in crime: POPX2 phosphatase and its interacting proteins in cancer. *Cell Death Dis* 2020; 11: 840. <https://doi.org/10.1038/s41419-020-03061-0>
- [39] JURMEISTER S, BAUMANN M, BALWIERZ A, KEKLIKOGLOU I, WARD A et al. MicroRNA-200c represses migration and invasion of breast cancer cells by targeting actin-regulatory proteins FHOD1 and PPM1F. *Mol Cell Biol* 2012; 32: 633-651. <https://doi.org/10.1128/MCB.06212-11>

Figure Legends

Figure 1. CircSEC24A was overexpressed in HCC. A) Venn diagram showed the overlapping circRNAs among the top 50 upregulated circRNAs in the GSE78520 and GSE94508 microarray datasets. B) Schematic illustration presented the formation of circSEC24A. C) RT-qPCR analysis of circSEC24A expression in 40 paired HCC tissues and peri-tumor tissues. D) Kaplan-Meier survival

447 curve and log-rank test were employed to assess the overall survival probability of HCC patients
448 with high and low circSEC24A expression. Low and high circSEC24A expression was cut off by
449 the median expression. E) Expression level of circSEC24A in HCC cells (Huh7, SK-Hep-1, Hep3B,
450 and Bel-7405) and THEL-2 cells was detected by RT-qPCR. F) The circular structure of
451 circSEC24A was verified by RNase R treatment and RT-qPCR analysis. *p < 0.05

452

453 **Figure 2.** CircSEC24A facilitated malignant behaviors of HCC cells. A) RT-qPCR verified the
454 knockdown efficiency of si-circSEC24A#1, si-circSEC24A#2, and si-circSEC24A#3 in Huh7 and
455 Hep3B cells. B-H) Influence of circSEC24A inhibition on growth, colony formation, cell cycle
456 progression, and proliferation of Huh7 and Hep3B cells was analyzed by CCK-8, colony formation,
457 flow cytometry, and EDU assays. I-N) Impacts of circSEC24A silencing on apoptosis, metastasis,
458 and invasion of Huh7 and Hep3B cells were assessed by flow cytometry and transwell assays. O, P)
459 Protein levels of E-cadherin and Vimentin in circSEC24A-repressed Huh7 and Hep3B cells were
460 measured by western blotting. *p < 0.05

461

462 **Figure 3.** CircSEC24A served as a sponge for miR-455-3p. A) Schematic illustration presented the
463 binding sites between circSEC24A and miR-455-3p. B) Dual-luciferase reporter detected the
464 luciferase activity of the circSEC24A wt or circSEC24A mut reporter in 293T cells transfected with
465 miR-455-3p mimic and miR-NC. C) RIP was verified to verify the interaction between circSEC24A
466 and miR-455-3p. D, E) RT-qPCR detected the expression of miR-455-3p in HCC tissues and cells.
467 F) Pearson's correlation analysis revealed the correlation between circSEC24A and miR-455-3p in
468 HCC tissues. *p < 0.05

469

470 **Figure 4.** CircSEC24A adsorbed miR-455-3p to accelerate proliferation, metastasis, invasion, and
471 EMT of HCC cells. A) RT-qPCR assessed the expression of miR-455-3p in Huh7 and Hep3B cells
472 transfected with miR-455-3p inhibitor or inhibitor NC. B-M) Huh7 and Hep3B cells were
473 transfected with si-NC, si-circSEC24A#1, si-circSEC24A#1+inhibitor NC, or
474 si-circSEC24A#1+miR-455-3p inhibitor. (B) Analysis of the influence of miR-455-3p inhibitor on
475 the expression of miR-455-3p in si-circSEC24A#1-transfected Huh7 and Hep3B cells by RT-qPCR.
476 (C-F) Assessment of the effect of miR-455-3p inhibitor on growth, colony formation, cell cycle

477 progression, and proliferation of si-circSEC24A#1-transfected Huh7 and Hep3B cells by CCK-8,
 478 colony formation, flow cytometry, and EDU assays. G-K) Impacts of miR-455-3p inhibitor on
 479 apoptosis, metastasis, and invasion of si-circSEC24A#1-transfected Huh7 and Hep3B cells were
 480 analyzed by flow cytometry and transwell assays. L, M) Western blotting assessed the influence of
 481 miR-455-3p inhibitor on protein levels of E-cadherin and Vimentin in si-circSEC24A#1-transfected
 482 Huh7 and Hep3B cells. * $p < 0.05$

484 **Figure 5.** CircSEC24A sponged miR-455-3p to regulate PPM1F expression. A) The predicted
 485 binding sites between miR-455-3p and PPM1F 3'UTR. B) Dual-luciferase reporter assay was
 486 carried out to detect the luciferase activity in 293T cells co-transfected with miR-455-3p mimic or
 487 miR-NC and the PPM1F 3'UTR wt or PPM1F 3'UTR mut reporter C, D) The levels of PPM1F
 488 mRNA and protein in HCC tissues and peri-tumor tissues were analyzed by RT-qPCR or western
 489 blotting. E) Protein level of PPM1F in HCC cells was assessed by western blotting. F) The
 490 correlation between PPM1F mRNA and miR-455-3p expression in HCC was determined by
 491 Pearson's correlation analysis. G) Influence of miR-455-3p inhibitor on the level of PPM1F protein
 492 in si-circSEC24A#1-transfected Huh7 and Hep3B cells was analyzed by western blotting. * $p < 0.05$

494 **Figure 6.** MiR-455-3p constrained HCC cell proliferation, metastasis, invasion, and EMT via
 495 targeting PPMF1. A) RT-qPCR validated the overexpression efficiency of miR-455-3p mimic in
 496 Huh7 and Hep3B cells. B) Western blotting confirmed the overexpression efficiency of
 497 pcDNA-PPMF1 in Huh7 and Hep3B cells. C-N) Huh7 and Hep3B cells were transfected with
 498 mimic NC, miR-455-3p mimic, miR-455-3p mimic+pcDNA, or miR-455-3p
 499 mimic+pcDNA-PPMF1. C-G) Analysis of the influence of PPMF1 overexpression on growth,
 500 colony formation, cell cycle progression, and proliferation of Huh7 and Hep3B cells carrying
 501 miR-455-3p mimic by CCK-8, colony formation, flow cytometry, and EDU assays. H-L)
 502 Assessment of the effect of PPMF1 overexpression on apoptosis, metastasis, and invasion of Huh7
 503 and Hep3B cells transfected with si-circSEC24A#1 by flow cytometry and transwell assays. M, N)
 504 Analysis of the effect of PPMF1 overexpression on protein levels of E-cadherin and Vimentin in
 505 si-circSEC24A#1-transfected Huh7 and Hep3B cells by western blotting. * $p < 0.05$

507 **Figure 7.** CircSEC24A silencing curbed HCC growth *in vivo*. A) Tumor volume was recorded once
508 a week after mice injection with stable Hep3B cells. B) Average tumor weight was calculated after
509 xenograft tumor removal. C) RT-qPCR was carried out to assess the expression levels of
510 circSEC24A, miR-455-3p, and PPM1F mRNA in xenograft tumors. D) IHC analysis revealed
511 protein levels of PPM1F, E-cadherin, and Vimentin in xenograft tumors. * $p < 0.05$
512

Accepted manuscript

513 **Table 1.** Primer sequences used for RT-qPCR analysis.

Genes	Primer sequences (5'-3')
circSEC24A	Forward (F): 5'-GCCTGGACTCATGGTTCCTT-3' Reverse (R): 5'-TGCTGACCAGAACAGTCCAA-3'
SEC24A	F: 5'-AGCCCTGAAACACGAGAGGAA-3' R: 5'-TGCTGACCAGAACAGTCCAAGG-3'
PPM1F	F: 5'-GCTGCTACAGACAGACCTTTCC-3' R: 5'-GGCGGTTAAAGAAACTCTGTGCC-3'
GAPDH	F: 5'-GTCTCCTCTGACTTCAACAGCG-3' R: 5'-ACCACCTGTGCTGTAGCCAA-3'
miR-455-3p	F: 5'-CGGCAGTCCATGGGCAT-3' R: 5'-AGTGCAGGGTCCGAGGTATT-3'
U6	F: 5'-CTCGCTTCGGCAGCACA-3' R: 5'-AACGCTTCACGAATTTGCGT-3'

514

Fig. 1 [Download full resolution image](#)

Fig. 2 [Download full resolution image](#)

Fig. 3 [Download full resolution image](#)

Fig. 4 [Download full resolution image](#)

Fig. 5 [Download full resolution image](#)

Fig. 6 [Download full resolution image](#)

Fig. 7 [Download full resolution image](#)